

# Vajra™

Fipronil 15% + Deltamethrin 2.5% w/v SC  
(Insecticide)

## Read Leaflet Before Use.

Fipronil 15% + Deltamethrin 2.5% w/v SC is a suspension concentrate containing Fipronil 15% (a.i.) and Deltamethrin 2.5% (a.i.) in the formulation. It is used for the control of tomato fruit borer on Tomato.

Crop	Common Name of the Pest	Dosage/HA		Dilution in water (liter)	Pre harvest interval (Days)	Re-entry Period (Days)
		g/L	Formulation (ml)			
Tomato	Tomato fruit borer	75 12.5	500	500	7	48

**Shake well before use**  
Good Practices relating to pesticide application techniques  
**Direction of use:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Weather condition (in general):** Spray preferably during morning and evening hours. Do not apply on a windy day or after rain.  
**Soil and water (where ever applicable):** Only use as soil application.

**Crop stage for Application:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity.

**Application Techniques:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Equipment Type:** Knapsack sprayer

**Notice:** Do not use above recommended tank for dosage details.

**Instruction for Mixing:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Re-entry Period:** 48 hours.

**Notice & Time of application:** Avoid application in afternoon hours. Ideal timing for application is morning and evening hours. Do not spray against wind.

**Application Timings:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity. Do not spray on a windy day or during rain.

**Frequency including maximum number of sprays:** Apply first spray during initial appearance of pests and repeat once or two sprays at 10 days interval depending upon pest intensity. Maximum two sprays.

**Stages during which pesticides should not be applied:** on any crop. Apply only in the stages as indicated on the label. Do not apply on any other stage.

**Symptoms of Poisoning:** Nausea/vomiting, anxiety, muscle tremors, nausea, vomiting, dizziness, diarrhoea, and in extreme cases possibly convulsions may occur.

**First Aid Measures:** Call the doctor immediately. **If Ingested:** If person is conscious and alert, rinse mouth with water and give 1 glass of water to drink. Never give anything by mouth to an unconscious person. **On the skin:** Do not use undiluted material and unless advised by attending physician, if in contact. Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration. Consult a physician. In case of eye contact, wash thoroughly the affected part of body with soap and plenty of water including hands and under fingernails. If pain, redness, inflammation, or conjunctivitis develops, consult a physician. Wash segregated and thoroughly washed with soap and water separately before use. In case of eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, redness, inflammation, or conjunctivitis persists, consult eye specialist.

**Antidote:** No specific antidote is known. Treat symptomatically. Antihistamines may be given to control allergies.

**Toxicology:** Non-toxic to honeybees at recommended dose

**Storage Conditions:** 1. Storage and transportation of the product should be done in such a way that no foodstuff and animal feed are contaminated. 2. Store in a cool, dry well-ventilated place. 3. Do not store, keep, eat and drink anything from the original containers, away from foodstuff and animal feeds. 4. If leakage or spillage occurs during transportation or storage, necessary measures should be taken to prevent poisoning and pollution of soil and water. Keep out of reach of children.

**Safety Precautions:** 1. Keep away from foodstuffs, empty foodstuff containers and animal feed. 2. Avoid contact with mouth, eyes and skin. 3. Avoid inhaling the spray mist. 4. Wash thoroughly the contaminated faces and parts of the body after spraying. 5. Do not smoke, eat and drink anything while spraying. 6. Wear full protective clothing while mixing and spraying in the field. 7. Do not apply in afternoon hours. Ideal timing for application is morning & evening hours. 8. Do not spray against wind. 9. Spray only during rain or when there is no rain or not predicted for the next 12 hrs. 10. Do not spray treated crops in the field.

**Cautions:** 1. Not to be used for any purpose other than those specified on label and leaflets. 2. Avoid the container after use as directed on label. The product is highly toxic to fish, and honey bees, and thus product should be avoided to use in drainage water bodies and during the active growing period of bees. 3. Do not apply more than recommended dose per ha per application. 4. Do not use in fields under irrigation and in areas where it is not practiced. Spray this product in calm weather when there is no rain or no predicted rain for the next 12 hours. 5. Do not use in areas where spraying equipment/sprayers over the treated area to avoid runoff to water bodies or drainage systems. Do not spray directly to surface water. Water bodies and water courses should be kept clear of spray. Do not use during or after application.

**Disposal of used containers, washing from equipment and surplus product:** 1. Empty the containers. 2. Wash the containers, equipment and surplus packages to dispose packages or surplus material and washings in a safe manner. 3. Do not use for any other purpose. 4. Spray this product in sealed packages should not left outside to prevent re-use. 5. The packages after use be broken and buried away from habitation.

Chemical Composition:		
Fipronil 15%	15.0%	w/v
Deltamethrin 2.5%	2.5%	w/v
Sunflower oil	5.0%	w/v
Amino salt of Polyoxynolone alcohol and other Phosphate	4.0%	w/v
Alkyl Polyglycidyl Ether block Copolymer	4.0%	w/v
Dithiopyr Chlor	0.5%	w/v
Xanthum Gum	0.5%	w/v
20% aqueous diisopropyl ester solution of 1,2-dichlorobenzene	0.5%	w/v
Silicon emulsion diameter=	1.0%	w/v
Deionized water	Q.S.	
<b>Total</b>	<b>100.00%</b>	<b>W/W</b>

**Density:** 1.06 - 1.12 " DANGEROUS TO RE USE EMPTY CONTAINER "

Manufactured & Marketed by: GHARDA CHEMICALS LIMITED  
Regd. Office: Gharda House, 48 Hill Road, Bandra (W), Mumbai - 400 050, Factory: Industrial Growth Centre, Phase II, Samba, Dist. Jammu, Pin 184 121, Jammu & Kashmir

फिप्रोनिल १५% + डेल्टामेथिन २.५% w/v आ एस.सी.  
(कीटनाशक)

## इस्तेमाल से पहले संलग्न पत्रिका पढ़ें।

Fipronil 15% + Deltamethrin 2.5% w/v SC is a suspension concentrate containing Fipronil 15% (a.i.) and Deltamethrin 2.5% (a.i.) in the formulation. It is used for the control of tomato fruit borer on Tomato.

Crop	Common Name of the Pest	Dosage/HA		Dilution in water (liter)	Pre harvest interval (Days)	Re-entry Period (Days)
		g/L	Formulation (ml)			
Tomato	Tomato fruit borer	75 12.5	500	500	7	48

**Shake well before use**  
Good Practices relating to pesticide application techniques  
**Direction of use:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Weather condition (in general):** Spray preferably during morning and evening hours. Do not apply on a windy day or after rain.  
**Soil and water (where ever applicable):** Only use as soil application.

**Crop stage for Application:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity.

**Application Techniques:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Equipment Type:** Knapsack sprayer

**Notice:** Do not use above recommended tank for dosage details.

**Instruction for Mixing:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Re-entry Period:** 48 hours.

**Notice & Time of application:** Avoid application in afternoon hours. Ideal timing for application is morning and evening hours. Do not spray against wind.

**Application Timings:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity. Do not spray on a windy day or during rain.

**Frequency including maximum number of sprays:** Apply first spray during initial appearance of pests and repeat once or two sprays at 10 days interval depending upon pest intensity. Maximum two sprays.

**Stages during which pesticides should not be applied:** on any crop. Apply only in the stages as indicated on the label. Do not apply on any other stage.

**Symptoms of Poisoning:** Nausea/vomiting, anxiety, muscle tremors, nausea, vomiting, dizziness, diarrhoea, and in extreme cases possibly convulsions may occur.

**First Aid Measures:** Call the doctor immediately. **If Ingested:** If person is conscious and alert, rinse mouth with water and give 1 glass of water to drink. Never give anything by mouth to an unconscious person. **On the skin:** Do not use undiluted material and unless advised by attending physician, if in contact. Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration. Consult a physician. In case of eye contact, wash thoroughly the affected part of body with soap and plenty of water including hands and under fingernails. If pain, redness, inflammation, or conjunctivitis develops, consult a physician. Wash segregated and thoroughly washed with soap and water separately before use. In case of eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, redness, inflammation, or conjunctivitis persists, consult eye specialist.

**Antidote:** No specific antidote is known. Treat symptomatically. Antihistamines may be given to control allergies.

**Toxicology:** Non-toxic to honeybees at recommended dose

**Storage Conditions:** 1. Storage and transportation of the product should be done in such a way that no foodstuff and animal feed are contaminated. 2. Store in a cool, dry well-ventilated place. 3. Do not store, keep, eat and drink anything from the original containers, away from foodstuff and animal feeds. 4. If leakage or spillage occurs during transportation or storage, necessary measures should be taken to prevent poisoning and pollution of soil and water. Keep out of reach of children.

**Safety Precautions:** 1. Keep away from foodstuffs, empty foodstuff containers and animal feed. 2. Avoid contact with mouth, eyes and skin. 3. Avoid inhaling the spray mist. 4. Wash thoroughly the contaminated faces and parts of the body after spraying. 5. Do not smoke, eat and drink anything while spraying. 6. Wear full protective clothing while mixing and spraying in the field. 7. Do not apply in afternoon hours. Ideal timing for application is morning & evening hours. 8. Do not spray against wind. 9. Spray only during rain or when there is no rain or not predicted for the next 12 hrs. 10. Do not spray treated crops in the field.

**Cautions:** 1. Not to be used for any purpose other than those specified on label and leaflets. 2. Avoid the container after use as directed on label. The product is highly toxic to fish, and honey bees, and thus product should be avoided to use in drainage water bodies and during the active growing period of bees. 3. Do not apply more than recommended dose per ha per application. 4. Do not use in fields under irrigation and in areas where it is not practiced. Spray this product in calm weather when there is no rain or no predicted rain for the next 12 hours. 5. Do not use in areas where spraying equipment/sprayers over the treated area to avoid runoff to water bodies or drainage systems. Do not spray directly to surface water. Water bodies and water courses should be kept clear of spray. Do not use during or after application.

**Disposal of used containers, washing from equipment and surplus product:** 1. Empty the containers. 2. Wash the containers, equipment and surplus packages to dispose packages or surplus material and washings in a safe manner. 3. Do not use for any other purpose. 4. Spray this product in sealed packages should not left outside to prevent re-use. 5. The packages after use be broken and buried away from habitation.

Chemical Composition:		
Fipronil 15%	15.0%	w/v
Deltamethrin 2.5%	2.5%	w/v
Sunflower oil	5.0%	w/v
Amino salt of Polyoxynolone alcohol and other Phosphate	4.0%	w/v
Alkyl Polyglycidyl Ether block Copolymer	4.0%	w/v
Dithiopyr Chlor	0.5%	w/v
Xanthum Gum	0.5%	w/v
20% aqueous diisopropyl ester solution of 1,2-dichlorobenzene	0.5%	w/v
Silicon emulsion diameter=	1.0%	w/v
Deionized water	Q.S.	
<b>Total</b>	<b>100.00%</b>	<b>W/W</b>

**Density:** 1.06 - 1.12 " DANGEROUS TO RE USE EMPTY CONTAINER "

Manufactured & Marketed by: GHARDA CHEMICALS LIMITED  
Regd. Office: Gharda House, 48 Hill Road, Bandra (W), Mumbai - 400 050, Factory: Industrial Growth Centre, Phase II, Samba, Dist. Jammu, Pin 184 121, Jammu & Kashmir

फिप्रोनिल १५% + डेल्टामेथिन २.५% w/v आ एस.सी.  
(कीटनाशक)

## वापरण्याची माहिती पत्रका वाचा.

Fipronil 15% + Deltamethrin 2.5% w/v SC is a suspension concentrate containing Fipronil 15% (a.i.) and Deltamethrin 2.5% (a.i.) in the formulation. It is used for the control of tomato fruit borer on Tomato.

Crop	Common Name of the Pest	Dosage/HA		Dilution in water (liter)	Pre harvest interval (Days)	Re-entry Period (Days)
		g/L	Formulation (ml)			
Tomato	Tomato fruit borer	75 12.5	500	500	7	48

**Shake well before use**  
Good Practices relating to pesticide application techniques  
**Direction of use:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Weather condition (in general):** Spray preferably during morning and evening hours. Do not apply on a windy day or after rain.  
**Soil and water (where ever applicable):** Only use as soil application.

**Crop stage for Application:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity.

**Application Techniques:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Equipment Type:** Knapsack sprayer

**Notice:** Do not use above recommended tank for dosage details.

**Instruction for Mixing:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Re-entry Period:** 48 hours.

**Notice & Time of application:** Avoid application in afternoon hours. Ideal timing for application is morning and evening hours. Do not spray against wind.

**Application Timings:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity. Do not spray on a windy day or during rain.

**Frequency including maximum number of sprays:** Apply first spray during initial appearance of pests and repeat once or two sprays at 10 days interval depending upon pest intensity. Maximum two sprays.

**Stages during which pesticides should not be applied:** on any crop. Apply only in the stages as indicated on the label. Do not apply on any other stage.

**Symptoms of Poisoning:** Nausea/vomiting, anxiety, muscle tremors, nausea, vomiting, dizziness, diarrhoea, and in extreme cases possibly convulsions may occur.

**First Aid Measures:** Call the doctor immediately. **If Ingested:** If person is conscious and alert, rinse mouth with water and give 1 glass of water to drink. Never give anything by mouth to an unconscious person. **On the skin:** Do not use undiluted material and unless advised by attending physician, if in contact. Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration. Consult a physician. In case of eye contact, wash thoroughly the affected part of body with soap and plenty of water including hands and under fingernails. If pain, redness, inflammation, or conjunctivitis develops, consult a physician. Wash segregated and thoroughly washed with soap and water separately before use. In case of eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, redness, inflammation, or conjunctivitis persists, consult eye specialist.

**Antidote:** No specific antidote is known. Treat symptomatically. Antihistamines may be given to control allergies.

**Toxicology:** Non-toxic to honeybees at recommended dose

**Storage Conditions:** 1. Storage and transportation of the product should be done in such a way that no foodstuff and animal feed are contaminated. 2. Store in a cool, dry well-ventilated place. 3. Do not store, keep, eat and drink anything from the original containers, away from foodstuff and animal feeds. 4. If leakage or spillage occurs during transportation or storage, necessary measures should be taken to prevent poisoning and pollution of soil and water. Keep out of reach of children.

**Safety Precautions:** 1. Keep away from foodstuffs, empty foodstuff containers and animal feed. 2. Avoid contact with mouth, eyes and skin. 3. Avoid inhaling the spray mist. 4. Wash thoroughly the contaminated faces and parts of the body after spraying. 5. Do not smoke, eat and drink anything while spraying. 6. Wear full protective clothing while mixing and spraying in the field. 7. Do not apply in afternoon hours. Ideal timing for application is morning & evening hours. 8. Do not spray against wind. 9. Spray only during rain or when there is no rain or not predicted for the next 12 hrs. 10. Do not spray treated crops in the field.

**Cautions:** 1. Not to be used for any purpose other than those specified on label and leaflets. 2. Avoid the container after use as directed on label. The product is highly toxic to fish, and honey bees, and thus product should be avoided to use in drainage water bodies and during the active growing period of bees. 3. Do not apply more than recommended dose per ha per application. 4. Do not use in fields under irrigation and in areas where it is not practiced. Spray this product in calm weather when there is no rain or no predicted rain for the next 12 hours. 5. Do not use in areas where spraying equipment/sprayers over the treated area to avoid runoff to water bodies or drainage systems. Do not spray directly to surface water. Water bodies and water courses should be kept clear of spray. Do not use during or after application.

**Disposal of used containers, washing from equipment and surplus product:** 1. Empty the containers. 2. Wash the containers, equipment and surplus packages to dispose packages or surplus material and washings in a safe manner. 3. Do not use for any other purpose. 4. Spray this product in sealed packages should not left outside to prevent re-use. 5. The packages after use be broken and buried away from habitation.

Chemical Composition:		
Fipronil 15%	15.0%	w/v
Deltamethrin 2.5%	2.5%	w/v
Sunflower oil	5.0%	w/v
Amino salt of Polyoxynolone alcohol and other Phosphate	4.0%	w/v
Alkyl Polyglycidyl Ether block Copolymer	4.0%	w/v
Dithiopyr Chlor	0.5%	w/v
Xanthum Gum	0.5%	w/v
20% aqueous diisopropyl ester solution of 1,2-dichlorobenzene	0.5%	w/v
Silicon emulsion diameter=	1.0%	w/v
Deionized water	Q.S.	
<b>Total</b>	<b>100.00%</b>	<b>W/W</b>

**Density:** 1.06 - 1.12 " DANGEROUS TO RE USE EMPTY CONTAINER "

Manufactured & Marketed by: GHARDA CHEMICALS LIMITED  
Regd. Office: Gharda House, 48 Hill Road, Bandra (W), Mumbai - 400 050, Factory: Industrial Growth Centre, Phase II, Samba, Dist. Jammu, Pin 184 121, Jammu & Kashmir

ફિપ્રોનિલ 15% + ડેલ્ટામેથ્રિન 2.5% w/v આ એસ.સી.  
(કીટનાશક)

## વાપરતી પહેલાં પત્રિકા વાંચો.

Fipronil 15% + Deltamethrin 2.5% w/v SC is a suspension concentrate containing Fipronil 15% (a.i.) and Deltamethrin 2.5% (a.i.) in the formulation. It is used for the control of tomato fruit borer on Tomato.

Crop	Common Name of the Pest	Dosage/HA		Dilution in water (liter)	Pre harvest interval (Days)	Re-entry Period (Days)
		g/L	Formulation (ml)			
Tomato	Tomato fruit borer	75 12.5	500	500	7	48

**Shake well before use**  
Good Practices relating to pesticide application techniques  
**Direction of use:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Weather condition (in general):** Spray preferably during morning and evening hours. Do not apply on a windy day or after rain.  
**Soil and water (where ever applicable):** Only use as soil application.

**Crop stage for Application:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity.

**Application Techniques:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Equipment Type:** Knapsack sprayer

**Notice:** Do not use above recommended tank for dosage details.

**Instruction for Mixing:** Measure required quantity of the product and mix it well with a small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable knapsack sprayer fitted with hi-flow cone nozzle.

**Re-entry Period:** 48 hours.

**Notice & Time of application:** Avoid application in afternoon hours. Ideal timing for application is morning and evening hours. Do not spray against wind.

**Application Timings:** Apply first spray during initial appearance of pests and repeat once in two sprays at 10 days interval depending upon pest intensity. Do not spray on a windy day or during rain.

**Frequency including maximum number of sprays:** Apply first spray during initial appearance of pests and repeat once or two sprays at 10 days interval depending upon pest intensity. Maximum two sprays.

**Stages during which pesticides should not be applied:** on any crop. Apply only in the stages as indicated on the label. Do not apply on any other stage.

**Symptoms of Poisoning:** Nausea/vomiting, anxiety, muscle tremors, nausea, vomiting, dizziness, diarrhoea, and in extreme cases possibly convulsions may occur.

**First Aid Measures:** Call the doctor immediately. **If Ingested:** If person is conscious and alert, rinse mouth with water and give 1 glass of water to drink. Never give anything by mouth to an unconscious person. **On the skin:** Do not use undiluted material and unless advised by attending physician, if in contact. Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration. Consult a physician. In case of eye contact, wash thoroughly the affected part of body with soap and plenty of water including hands and under fingernails. If pain, redness, inflammation, or conjunctivitis develops, consult a physician. Wash segregated and thoroughly washed with soap and water separately before use. In case of eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, redness, inflammation, or conjunctivitis persists, consult eye specialist.

**Antidote:** No specific antidote is known. Treat symptomatically. Antihistamines may be given to control allergies.

**Toxicology:** Non-toxic to honeybees at recommended dose

**Storage Conditions:** 1. Storage and transportation of the product should be done in such a way that no foodstuff and animal feed are contaminated. 2. Store in a cool, dry well-ventilated place. 3. Do not store, keep, eat and drink anything from the original containers, away from foodstuff and animal feeds. 4. If leakage or spillage occurs during transportation or storage, necessary measures should be taken to prevent poisoning and pollution of soil and water. Keep out of reach of children.

**Safety Precautions:** 1. Keep away from foodstuffs, empty foodstuff containers and animal feed. 2. Avoid contact with mouth, eyes and skin. 3. Avoid inhaling the spray mist. 4. Wash thoroughly the contaminated faces and parts of the body after spraying. 5. Do not smoke, eat and drink anything while spraying. 6. Wear full protective clothing while mixing and spraying in the field. 7. Do not apply in afternoon hours. Ideal timing for application is morning & evening hours. 8. Do not spray against wind. 9. Spray only during rain or when there is no rain or not predicted for the next 12 hrs. 10. Do not spray treated crops in the field.

**Cautions:** 1. Not to be used for any purpose other than those specified on label and leaflets. 2. Avoid the container after use as directed on label. The product is highly toxic to fish, and honey bees, and thus product should be avoided to use in drainage water bodies and during the active growing period of bees. 3. Do not apply more than recommended dose per ha per application. 4. Do not use in fields under irrigation and in areas where it is not practiced. Spray this product in calm weather when there is no rain or no predicted rain for the next 12 hours. 5. Do not use in areas where spraying equipment/sprayers over the treated area to avoid runoff to water bodies or drainage systems. Do not spray directly to surface water. Water bodies and water courses should be kept clear of spray. Do not use during or after application.

**Disposal of used containers, washing from equipment and surplus product:** 1. Empty the containers. 2. Wash the containers, equipment and surplus packages to dispose packages or surplus material and washings in a safe manner. 3. Do not use for any other purpose. 4. Spray this product in sealed packages should not left outside to prevent re-use. 5. The packages after use be broken and buried away from habitation.

Chemical Composition:		
Fipronil 15%	15.0%	w/v
Deltamethrin 2.		

